

NEW SPECTRAL MULTIPLICITIES FOR MIXING TRANSFORMATIONS

ALEXANDRE I. DANILENKO

ABSTRACT. It is shown that if E is any subset of $\mathbb{N}$ such that either $1 \in E$ or $2 \in E$ then there is a *mixing* transformation whose set of spectral multiplicities is E .

0. INTRODUCTION

Let T be an invertible measure preserving transformation of a standard probability non-atomic space $(X, \mathfrak{B}, \mu)$. Denote by U_T the associated unitary Koopman operator on the space $L^2(X, \mu)$:

$$U_T f = f \circ T.$$

Let $\mathcal{M}(T)$ stand for the set of essential values of the spectral multiplicity function for $U_T \upharpoonright (L^2(X, \mu) \ominus \mathbb{C})$. A subset $M \subset \mathbb{N}$ is called *realizable* if $M = \mathcal{M}(T)$ for an ergodic transformation T . The *spectral multiplicity problem* can be stated as follows

(Pr1) which subsets of $\mathbb{N}$ are realizable?

It is one of the basic problems of the spectral theory of dynamical systems. Many papers are devoted to partial solutions of (Pr1): [Os], [Ro1], [Ro3], [G–Li], [KL], [Ag1], [Ry1], [Ag2], [Ag3], [Da3], [Ag4], [KaL], [Da5], [Ry5] (see also a recent survey [Le]). In particular, it was shown there that the following subsets are realizable:

- every subset containing 1,
- every subset containing 2,
- subsets $n \cdot E$, where $n > 0$ and E is an arbitrary subset containing 1.

Despite of this progress, (Pr1) remains unsolved in general. A transformation T is said to be

- *mixing* if $\mu(T^n A \cap B) \rightarrow \mu(A)\mu(B)$ as $n \rightarrow \infty$ for all subsets $A, B \in \mathfrak{B}$,
- *rigid* if $\mu(T^{n_k} A \cap B) \rightarrow \mu(A \cap B)$ for some subsequence $n_k \rightarrow \infty$ and all subsets $A, B \in \mathfrak{B}$,
- *partially rigid* if $\mu(T^{n_k} A \cap B) \geq \delta \mu(A \cap B)$ along some subsequence $n_k \rightarrow \infty$ for certain $\delta > 0$ and all subsets $A, B \in \mathfrak{B}$.

We note that the realizations in the aforementioned papers are either rigid or partially rigid. The present paper is devoted to a more subtle question:

(Pr2) for which subsets E of $\mathbb{N}$ there is a *mixing* transformation T such that $\mathcal{M}(T) = E$?

We then say that E admits a *mixing realization*. The information about mixing realizations is rather scarce. The first mixing transformation with a simple spectrum (i.e. a mixing realization for $\{1\}$) was constructed in [Ne] as a Gaussian system. In [Ro2] it was shown that for each $n > 0$, there is a mixing T with $n < \sup \mathcal{M}(T) < \infty$. The paper [Ry3] provides mixing realizations for $\{2\}$. More generally, for each $n > 1$, Ageev builds mixing realizations for $\{2, 3, \dots, n\}$ in [Ag4]. It should be noted that his result follows also from [Ry4], where Ryzhikov constructs mixing transformations whose symmetric powers have all a simple spectrum. (We also mention that (P1) and (P2) are solved completely in the category of *infinite* measure preserving transformations [DaR1], [DaR2]).

The set of mixing transformations $\mathbb{M}$ is meager in the (Polish) weak topology on the group of all measure preserving transformations. Tikhonov [Ti] introduced a Polish topology on $\mathbb{M}$ which is strictly stronger than the induced weak topology. He showed, in particular, that a generic mixing transformation has a simple spectrum. Thus we are interested in realizations belonging to a meager part of $\mathbb{M}$. This somehow explains a complexity of (Pr2) comparatively with (Pr1): meager sets in Tikhonov's topology are “much smaller” than meager sets in the weak topology.

We now state the main result of the paper.

Main Theorem. *Given an arbitrary subset E of $\mathbb{N}$ such that either $1 \in E$ or $2 \in E$, there exists a mixing (of all orders) transformation S such that $\mathcal{M}(S) = E$.*

To prove the theorem we combine various techniques and approaches to spectral realizations developed by a number of authors:

- non-Abelian compact group extensions [Ro1]–[Ro3],
- natural factors of Abelian compact group extensions [KL], [G–L],
- spectral multiplicities of Cartesian products and their natural factors [Ka], [Ry1], [Ag1], [Ag4],
- weak limits of powers of the associated Koopman operator [Ag1], [Ry1], [KaL], etc.,
- techniques to force mixing [Ry4],
- mixing properties of staircase algorithms [Ad],
- (C, F) -construction [dJ], [Da2]–[Da5].

The outline of the paper is as follows. In Section 1 we consider cocycles of dynamical systems with values in compact group extensions and discuss their simplest spectral properties. (C, F) -construction of dynamical systems, an algebraic version of the classical cutting-and-stacking is outlined briefly in Section 2.

In Section 3 we prove a part of Main Theorem when $1 \in E$. It is worthy to note that the realizations from [G–Li] and [KL] based on Abelian compact group extensions are—on our opinion—inherently rigid and hence do not suit for this purpose. That is why we “return” to historically earlier non-Abelian group extensions that go back to [Ro1], [Ro3] (see also [KaL]). Since they were elaborated for spectral realization of only *finite* sets, we first show how to adjust them to *arbitrary* subsets of $\mathbb{N}$ via a “compactification” of the Algebraic Lemma from [KL]. Then we adapt Ryzhikov's idea *to force mixing* in symmetric products ([Ry3] and [Ry4]) to compact extensions. The idea is as follows. A (C, F) -construction of the extensions is implemented in such a way to ensure availability of some special weak limits in the closure of powers of the associated Koopman operators. These limits imply simple spectrum of some invariant components of the Koopman operators. That, in turn,

guarantees that the spectral multiplicities can be computed from a certain algebraic picture. Moreover, for each $\delta > 0$, it is possible to modify this construction in such a way to obtain a “mixing component” that “occupies” δ -part of the whole space. At the same time the “rigid component” occupying $(1 - \delta)$ -part of the space is retained to guarantee the necessary weak limits of powers of the Koopman operator. Then we *pass to a limit* along a sequence of such δ_i -“partly mixing” transformations which have the same (desired) spectral multiplicities as $\delta_i \rightarrow 1$. The limit will be a mixing compact extension of a rank-one mixing map. Though there are no non-trivial weak limits in the closure of powers of the limit Koopman operator any more, it is possible to retain the *simple spectrum* properties of the components of the Koopman operators in the limit. Hence the limit Koopman operator has the same spectral multiplicities as the prelimit ones.

Section 4 is devoted entirely to the proof of the remaining part of Main Theorem, i.e. when $2 \in E$. The scheme of the proof is as in the previous section. We note that the realizations (of all sets containing 2) constructed in [Da5] are immanently rigid. They are not suit to force mixing. Therefore we again use non-Abelian group extensions. For that we first generalize the Algebraic Lemma from [KL] in such a way to *embrace* the case when $1 \notin E$. It should be noted that our generalization implies the affirmative answer to the following question from [KaL]:

whether the series of weak mixing realizations constructed in [KaL] covers all finite subsets of $\mathbb{N}$ that contain 2?

Several concluding remarks are contained in Section 5.

The author thanks V. Ryzhikov for a suggestion to apply the techniques “to force mixing” from [Ry3] and [Ry4] to compact group extensions.

1. COCYCLES WITH VALES IN GROUP EXTENSIONS

Let T be an ergodic transformation of (X, μ) . Denote by $\mathcal{R} \subset X \times X$ the T -orbit equivalence relation. A Borel map α from $\mathcal{R}$ to a compact group K is called a *cocycle* of $\mathcal{R}$ if

$$\alpha(x, y)\alpha(y, z) = \alpha(x, z)$$

for all $(x, y), (y, z) \in \mathcal{R}$. Two cocycles $\alpha, \beta : \mathcal{R} \rightarrow K$ are *cohomologous* if

$$\alpha(x, y) = \phi(x)\beta(x, y)\phi(y)^{-1}$$

at a.a. $(x, y) \in \mathcal{R}$ for a Borel map $\phi : X \rightarrow K$. Denote by $C(T)$ the *centralizer* of T , i.e. the group of all transformations commuting with T . Given $S \in C(T)$, a cocycle $\alpha \circ S : \mathcal{R} \rightarrow K$ is well defined by

$$\alpha \circ S(x, y) := \alpha(Sx, Sy).$$

for all $(x, y) \in \mathcal{R}$. Let λ_K denote Haar measure on K . Define a transformation T_α of the probability space $(X \times K, \mu \times \lambda_K)$ by setting

$$T_\alpha(x, k) := (Tx, k\alpha(x, Tx)).$$

It is called the *α -skew product extension* of T . If it is ergodic then α is called *ergodic*. We note that the centralizer of T_α contains a compact subgroup K which acts on $X \times K$ by left translations along the second coordinate. We note that if

K is Abelian then U_{T_α} is unitarily equivalent to the orthogonal sum $\bigoplus_{\chi \in \widehat{K}} U_{T,\chi}$, where $U_{T,\chi}$ is a unitary operator in $L^2(X, \mu)$ given by

$$U_{T,\chi}f(x) = \chi(\alpha(x, Tx))f(Tx).$$

If H is a closed subgroup of K then we denote by $T_{\alpha,H}$ the following transformation of $(X \times H \setminus K, \mu \times \lambda_{H \setminus K})$:

$$T_{\alpha,H}(x, Hk) := (Tx, Hk\alpha(x, Tx)).$$

It is easy to see that $T_{\alpha,H}$ is a factor of T_α . It is called a *natural* factor of T_α . Notice that $U_{T_{\alpha,H}}$ is unitarily equivalent to $\bigoplus_{\chi \in \widehat{K/H}} U_{T,\chi}$.

Let A be a compact K -module. This means that A is a compact second countable Abelian group and K acts on A by group automorphisms. Denote by G the semidirect product $K \ltimes A$. Recall that the multiplication in G is given by

$$(k, a)(k', a') := (kk', a + k \cdot a').$$

Given a cocycle $\gamma : \mathcal{R} \rightarrow G$, we can write it as a pair of “coordinate” maps $\gamma = (\beta, \alpha)$. Then β is a cocycle of $\mathcal{R}$ with values in K . As for the map $\alpha : \mathcal{R} \rightarrow A$, it is not a cocycle but it satisfies the following equation

$$\alpha(x, y) + \beta(x, y) \cdot \alpha(y, z) = \alpha(x, z)$$

for all $(x, y), (y, z) \in \mathcal{R}$. Denote by $\mathcal{R}(\beta)$ the T_β -orbit equivalence relation on $X \times K$. It is easy to verify that

$$\mathcal{R}(\beta) = \{((x, k), (x', k\beta(x, x'))) \mid (x, x') \in \mathcal{R}, k \in K\}.$$

Consider a map

$$\tilde{\alpha} : \mathcal{R}(\beta) \ni ((x, k), (x', k\beta(x, x'))) \mapsto k \cdot \alpha(x, x') \in A.$$

Then $\tilde{\alpha}$ is a cocycle of $\mathcal{R}(\beta)$. Moreover, the γ -skew product extension T_γ coincides with the “double” extension $(T_\beta)_{\tilde{\alpha}}$ of T . Given $k \in K$, denote by L_k the following transformation of $(X \times K, \mu \times \lambda_K)$: $L_k(x, r) := (x, kr)$. Then $L_k \in C(T_\beta)$ and

$$\tilde{\alpha} \circ L_k = k \cdot \tilde{\alpha}.$$

This implies the following:

$$(1-1) \quad U_{T_{\beta,\chi}} \text{ is unitarily equivalent to } U_{T_{\beta,\chi \circ k}} \text{ for each } k \in K.$$

For more information about cocycles with values in locally compact group extensions we refer to [Da1] and references therein.

2. (C, F) -SYSTEMS AND (C, F) -COCYCLES

We recall the (C, F) -construction (see [dJ], [Da2]–[Da4]). Let two sequences $(C_n)_{n>0}$ and $(F_n)_{n\geq 0}$ of finite subsets in $\mathbb{Z}$ are given such that:

- $F_n = \{0, 1, \dots, h_n - 1\}$,
- $0 \in C_n$, $\#C_n > 1$,
- $F_n + C_{n+1} \subset F_{n+1}$,
- $(F_n + c) \cap (F_n + c') = \emptyset$ if $c \neq c'$, $c, c' \in C_{n+1}$,
- $\lim_{n \rightarrow \infty} h_n / (\#C_1 \cdots \#C_n) < \infty$.

Let $X_n := F_n \times C_{n+1} \times C_{n+2} \times \cdots$. Endow this set with the (compact Polish) product topology. The following map

$$(f_n, c_{n+1}, c_{n+2}, \dots) \mapsto (f_n + c_{n+1}, c_{n+2}, \dots)$$

is a topological embedding of X_n into X_{n+1} . We now set $X := \bigcup_{n \geq 0} X_n$ and endow it with the (locally compact Polish) inductive limit topology. Given $A \subset F_n$, we denote by $[A]_n$ the following cylinder: $\{x = (f, c_{n+1}, \dots) \in X_n \mid f \in A\}$. Then $\{[A]_n \mid A \subset F_n, n > 0\}$ is the family of all compact open subsets in X . It forms a base of the topology on X .

Let $\mathcal{R}$ stand for the *tail* equivalence relation on X : two points $x, x' \in X$ are $\mathcal{R}$ -equivalent if there is $n > 0$ such that $x = (f_n, c_{n+1}, \dots)$, $x' = (f'_n, c'_{n+1}, \dots) \in X_n$ and $c_m = c'_m$ for all $m > n$. There is only one probability (non-atomic) Borel measure μ on X which is invariant (and ergodic) under $\mathcal{R}$.

Now we define a transformation T of (X, μ) by setting

$$T(f_n, c_{n+1}, \dots) := (1 + f_n, c_{n+1}, \dots) \text{ whenever } f_n < h_n - 1, n > 0.$$

This formula defines T partly on X_n . When $n \rightarrow \infty$, T extends to the entire X (modulo a countable subset) as a μ -preserving invertible transformation. Moreover, the T -orbit equivalence relation coincides with $\mathcal{R}$ on the subset where T is defined. We call (X, μ, T) the (C, F) -dynamical system associated with $(C_{n+1}, F_n)_{n \geq 0}$.

From now on we will assume that $\#C_n \rightarrow \infty$ as $n \rightarrow \infty$.

If for each $n > 0$, there are $i_n \geq 0$ and $c_n \in C_n$ such that $c_n + ih_n + \frac{i(i-1)}{2} \in C_n$ for each $i = 1, \dots, i_n$ and $i_n / \#C_n \rightarrow 1$ as $n \rightarrow \infty$ then the corresponding (C, F) -system is called an *almost staircase* [Ry4]. It follows from [Ad] that if $\frac{i_n^2}{h_n} \rightarrow 0$ then the corresponding almost staircase transformation is mixing.

We recall a concept of (C, F) -cocycle (see [Da3]). Let G be a compact second countable group. Given a sequence of maps $\gamma_n : C_n \rightarrow G$, $n = 1, 2, \dots$, we first define a Borel cocycle $\gamma : \mathcal{R} \cap (X_0 \times X_0) \rightarrow G$ by setting

$$(2-1) \quad \gamma(x, x') := \gamma_1(c_1) \cdots \gamma_n(c_n) \gamma_n(c'_n)^{-1} \cdots \gamma_1(c'_1)^{-1},$$

if $x = (f_0, c_1, c_2, \dots) \in X_0$, $x' = (f'_0, c'_1, c'_2, \dots) \in X_0$ and $c_i = c'_i$ for all $i > n$. To extend γ to the entire $\mathcal{R}$, we first define a map $\pi : X \rightarrow X_0$ as follows. Given $x \in X$, let n be the least positive integer such that $x \in X_n$. Then $x = (f_n, c_{n+1}, \dots) \in X_n$. We set

$$\pi(x) := (\underbrace{0, \dots, 0}_{n+1 \text{ times}}, c_{n+1}, c_{n+2}, \dots) \in X_0.$$

Of course, $(x, \pi(x)) \in \mathcal{R}$. Now for each pair $(x, y) \in \mathcal{R}$, we let

$$\gamma(x, y) := \gamma(\pi(x), \pi(y)).$$

It is easy to verify that γ is a well defined cocycle of $\mathcal{R}$ with values in K . We call it *the (C, F) -cocycle associated with $(\gamma_n)_{n=1}^\infty$* .

Suppose now that $G = K \ltimes A$. Then we can write $\gamma = (\beta, \alpha)$ as above. Also, $\gamma_n = (\beta_n, \alpha_n)$ for some maps $\beta_n : C_n \rightarrow K$ and $\alpha_n : C_n \rightarrow A$. Then β is the (C, F) -cocycle associated with $(\beta_n)_{n>0}$. As for the mapping α , we deduce from (2-1) that

$$(2-2) \quad \begin{aligned} \alpha(x, x') &= \alpha_1(c_1) + \sum_{j=2}^n \beta_1(c_1) \cdots \beta_{j-1}(c_{j-1}) \cdot \alpha_j(c_j) \\ &\quad - \beta(x, x') \cdot \left[\alpha_1(c'_1) + \sum_{j=2}^n \beta_1(c'_1) \cdots \beta_{j-1}(c'_{j-1}) \cdot \alpha_j(c'_j) \right] \end{aligned}$$

for all $x = (f, c_1, c_2, \dots)$, $x' = (f', c'_1, c'_2, \dots) \in X_0$ such that $c_i = c'_i$ whenever $i > n$.

3. MIXING TRANSFORMATIONS WHOSE SET OF SPECTRAL MULTIPLICITIES CONTAINS 1

Let E be an arbitrary subset of $\mathbb{N}$ such that $1 \in E$. Our main purpose in this section is to prove the following theorem.

Theorem 3.1. *There exists a mixing (of all orders) transformation S such that $\mathcal{M}(S) = E$.*

Let K be a compact Abelian group, B a countable K -module and D a subgroup of B . We denote

$$L(K, B, D) := \{\#(\{k \cdot d \mid k \in K\} \cap D), d \in D \setminus \{0\}\}.$$

The lemma below plays an important role in the proof of Theorem 3.1. We show how to deduce it from the Algebraic Lemma from [KL].

Lemma 3.2. *There are a compact Abelian group K , a compact K -module A and a closed subgroup $H \subset A$ such that $L(K, \widehat{A}, \widehat{A/H}) = E$ and*

- (i) *K is the inverse limit of a sequence of finite Abelian groups $K_1 \leftarrow K_2 \leftarrow \cdots$, where the arrows are onto homomorphisms,*
- (ii) *A is the inverse limit of a sequence of finite K -modules $A_1 \leftarrow A_2 \leftarrow \cdots$, where the arrows are K -equivariant onto homomorphisms (more precisely, A_n is a K_n -module, i.e. $k \cdot a = k' \cdot a$ if $\pi_n(k) = \pi_n(k')$, where $\pi_n : K \rightarrow K_n$ stands for the canonical projection),*
- (iii) *there is a dense countable sub- K -module $A_0 \subset A$ such that the K -orbit of each $a \in A_0$ is finite.*

Proof. It is shown in [KL] that there is a compact group $A = \mathbb{T}^{k_1} \times \mathbb{T}^{k_2} \times \cdots$, a closed subgroup H of A and a group automorphism $\sigma = \sigma_1 \times \sigma_2 \cdots$ of A such that $L(\mathbb{Z}, \widehat{A}, \widehat{A/H}) = E$, where σ_i is the automorphism of $\mathbb{T}^{k_i}$ generated by the cyclic

permutation of the coordinates and $\widehat{A}$ is considered as a $\mathbb{Z}$ -module with respect to the automorphism which is dual to σ . Our first observation is that the argument from [KL] works as well if we replace every copy of $\mathbb{T}$ in A with $\mathbb{Z}/2\mathbb{Z}$. Next, it is easy to see that the closure of the cyclic group generated by σ in the group of all group automorphisms of A equipped with the natural Polish topology is the inverse limit of the sequence

$$\mathbb{Z}/k_1\mathbb{Z} \leftarrow \mathbb{Z}/(k_1k_2\mathbb{Z}) \leftarrow \dots$$

equipped with the natural compact topology. Denote this inverse limit by K . Then $L(\mathbb{Z}, \widehat{A}, \widehat{A/H}) = L(K, \widehat{A}, \widehat{A/H})$. The properties (i)–(iii) are now obvious. $\square$

From now on till the end of the section (K, A, H) denote the groups whose existence is asserted in Lemma 3.2.

We first construct a partially rigid transformation S with $\mathcal{M}(S) = E$ and whose *staircase (i.e. mixing) part* is $(1 - \delta)$ -large. For that we will use skew product extensions of (C, F) -transformations constructed via (C, F) -cocycles.

Fix a countable dense subgroup $K_0 \subset K$. Let $0 < i_n \leq r_n$, $r_n \rightarrow \infty$, $i_n/r_n \rightarrow \delta$, as $n \rightarrow \infty$, where $0 < \delta < 1$. We now let

$$c_{n+1}(i) := \begin{cases} 0, & \text{if } i = 0, \\ c_{n+1}(i-1) + h_n, & \text{if } 0 < i < i_n, \\ c_{n+1}(i-1) + h_n + i - i_n, & \text{if } i_n \leq i < r_n. \end{cases}$$

We define C_{n+1} by setting

$$C_{n+1} := \{c_{n+1}(i) \mid 0 \leq i < r_n\}.$$

Let $A_0 \subset A$ be a countable submodule which is dense in A . Consider a partition $\mathbb{N} = \bigsqcup_{a \in A_0} \mathcal{N}_a \sqcup \bigsqcup_{k \in K_0} \mathcal{L}_k$ of $\mathbb{N}$ into infinite subsets $\mathcal{N}_a, \mathcal{L}_k$.

If $n+1 \in \mathcal{N}_a$, we define a mapping $\alpha_{n+1} : C_{n+1} \rightarrow A$ by

$$\alpha_{n+1}(c_{n+1}(i)) := \begin{cases} 0, & \text{if } i = 0, \\ \alpha_{n+1}(c_{n+1}(i-1)) - a, & \text{if } i = 1, \dots, i_n - 1, \\ \alpha_{n+1}(c_{n+1}(i-1)), & \text{if } i = i_n, \dots, r_n - 1. \end{cases}$$

We also let $\beta_{n+1} \equiv 1$ on C_{n+1} .

If $n+1 \in \mathcal{L}_k$, we let $\alpha_{n+1} \equiv 0$ on C_{n+1} . We also define $\beta_{n+1} : C_{n+1} \rightarrow K$ by setting

$$\beta_{n+1}(c_{n+1}(i)) := \begin{cases} 0, & \text{if } i = 0, \\ \beta_{n+1}(c_{n+1}(i-1)) - k, & \text{if } i = 1, \dots, i_n - 1, \\ \beta_{n+1}(c_{n+1}(i-1)), & \text{if } i = i_n, \dots, r_n - 1. \end{cases}$$

Let (X, μ, T) stand for the (C, F) -system associated with $(C_{n+1}, F_n)_{n \geq 0}$. Denote by $\mathcal{R}$ the tail equivalence relation on X and denote by $\gamma = (\beta, \alpha) : \mathcal{R} \rightarrow G$ the (C, F) -cocycle associated with $(\beta_n, \alpha_n)_{n \geq 0}$. Recall that given a character $\chi \in \widehat{A}$, $U_{T_\gamma, \chi}$ stand for the following unitary operator on $L^2(X \times K, \mu \times \lambda_K)$:

$$U_{T_\beta, \chi} v(x, k) := \chi(\tilde{\alpha}((x, k), (Tx, k\beta(x, Tx)))) v(Tx, k\beta(x, Tx)).$$

We also note that $U_{T_\beta, 1} := U_{T_\beta}$. Given $a \in A_0$, we let $\text{Orb}_K(a) := \{k \cdot a \mid k \in K\}$ and

$$l_\chi(a) := \frac{1}{\#\text{Orb}_K(a)} \sum_{b \in \text{Orb}_K(a)} \chi(b)$$

for $\chi \in \widehat{A}$ and $a \in A$.

Proposition 3.4. For each $a \in A_0$,

- (i) $U_{T_\beta, 1}^{h_n} \rightarrow \delta I + (1 - \delta)P_0$ weakly as $\mathcal{N}_a - 1 \ni n \rightarrow \infty$, where P_0 is the orthogonal projection onto the subspace of constant functions on $X \times K$.
- (ii) $U_{T_\beta, \chi}^{h_n} \rightarrow \delta l_\chi(a)I$ weakly as $\mathcal{N}_a - 1 \ni n \rightarrow \infty$ for each $1 \neq \chi \in \widehat{A}$.

For each $k \in K_0$,

- (iii) $U_{T, 1}^{h_n} \rightarrow \delta I + (1 - \delta)P_0$ weakly as $\mathcal{L}_k - 1 \ni n \rightarrow \infty$, where P_0 is the orthogonal projection onto the subspace of constant functions on X .
- (iv) $U_{T, \eta}^{h_n} \rightarrow \delta \eta(k)I$ weakly as $\mathcal{L}_k - 1 \ni n \rightarrow \infty$ for each $1 \neq \eta \in \widehat{K}$.

Proof. We first verify that β is ergodic. Take arbitrary $n > 0$, $k \in K$ and a subset $B \subset F_n$. Find $m > n$ such that $m \in \mathcal{L}_k - 1$. Let

$$I_m := \{0 \leq i < i_m - 1 \mid \}.$$

We now let

$$B_0 := B + C_{n+1} + \cdots + C_m + c_{m+1}(I_m) \subset F_{m+1}.$$

If $x = (f_m, c_{m+1}(i_1), c_{m+2}(i_2), \dots) \in X_m$ belongs to $[B_0]_{m+1}$ then $i_1 \in I_m$ and

$$T^{h_m}x = (f_m, c_{m+1}(i_1 + 1), c_{m+2}(i_2), \dots).$$

Moreover,

$$[B_0]_{m+1} \cup T^{h_m}[B_0]_{m+1} \subset [B]_n, \quad \mu([B_0]_{m+1}) > 0.5\delta\mu([B]_n),$$

and $\beta(x, T^{h_m}x) = k$ for all $x \in [B_0]_{m+1}$. By a standard criterium of ergodicity for cocycles (see [FM], [Sc]), β is ergodic.

Our next purpose is to show (i). Take $n \in \mathcal{N}_a - 1$, subsets $B, B' \subset F_n$ and $\eta \in \widehat{K}$. Then

$$\begin{aligned} (U_{T, \eta}^{h_n} 1_{[B]_n})(x) &= \eta(\beta(x, T^{h_n}x)) 1_{T^{-h_n}[B]_n}(x) \\ &= \sum_{j=1}^{i_n-1} \eta(\beta(x, T^{h_n}x)) 1_{[B+c_{n+1}(j-1)]_{n+1}}(x) \\ &\quad + \sum_{j=i_n+1}^{r_n-1} \eta(\beta(T^{i_n-j}x, T^{h_n}x)) U_{T_\beta, \eta}^{i_n-j} 1_{[B+c_{n+1}(j-1)]_{n+1}}(x) + \bar{o}(1) \\ &= \sum_{j=1}^{i_n-1} 1_{[B+c_{n+1}(j-1)]_{n+1}}(x) \\ &\quad + \sum_{j=i_n+1}^{r_n-1} U_{T, \eta}^{i_n-j} 1_{[B+c_{n+1}(j-1)]_{n+1}}(x) + \bar{o}(1) \end{aligned}$$

for all $x \in X$ and $k \in K$. Then

$$\langle U_{T, \eta}^{h_n} 1_{[B]_n}, 1_{[B']_n} \rangle = \frac{i_n}{r_n} \langle 1_{[B]_n}, 1_{[B']_n} \rangle + \left\langle \frac{1}{r_n} \sum_{j=1}^{r_n-i_n} U_{T, \eta}^{-j} 1_{[B]_n}, 1_{[B']_n} \right\rangle + \bar{o}(1).$$

It follows from the mean ergodic theorem that $(r_n - i_n)^{-1} \sum_{j=1}^{r_n - i_n} U_{T,\eta}^{-j}$ converges strongly to the orthogonal projection P_η onto the subspace of $U_{T,\eta}$ -invariant vectors. Hence

$$U_{T,\eta}^{h_n} \rightarrow \delta I + (1 - \delta)P_\eta \text{ weakly as } \mathcal{N}_a - 1 \ni n \rightarrow \infty.$$

If $\eta = 1$ then $P_\eta = P_0$ since T is ergodic. If $\eta \neq 1$ and $U_{T_\beta,\eta}v = v$ for a non-trivial function $v \in L^2(X, \mu)$ then $\eta(\beta(x, Tx))v(Tx) = v(x)$ at a.a. x . It follows that $\eta \circ \beta$ is cohomologous to the trivial cocycle. However this contradicts to the fact that β is ergodic. Hence $v = 0$ and therefore $P_\eta = 0$. Since $U_{T_\beta,1} = U_{T_\beta} = \bigoplus_{\eta \in \widehat{K}} U_{T,\eta}$, the claim (i) follows.

Claims (iii) and (iv) are shown in a similar way.

It remains to prove (ii). Fix $n \in \mathcal{N}_a - 1$. Take $B \subset F_n$, $\eta \in \widehat{K}$ and $\chi \in \widehat{A}$. It follows that

$$\begin{aligned} (3-1) \quad & U_{T_\beta,\chi}^{h_n}(1_{[B]_n} \otimes \eta)(x, k) = \chi(k \cdot \alpha(x, T^{h_n}x))1_{T^{-h_n}[B]_n}(x)\eta(k\beta(x, T^{h_n}x)) \\ & = \sum_{j=1}^{i_n-1} \chi(k \cdot \alpha(x, T^{h_n}x))1_{[B+c_{n+1}(j-1)]_{n+1}}(x)\eta(k) \\ & + \sum_{j=i_n+1}^{r_n-1} \eta(\beta(T^{i_n-j}x, T^{h_n}x))U_{T_\beta,\chi}^{i_n-j}(1_{[B+c_{n+1}(j-1)]_{n+1}} \otimes \eta)(x, k) + \bar{o}(1) \end{aligned}$$

for all $x \in X$ and $k \in K$.

We say that two points $x, y \in X$ are $\mathcal{R}_n$ -equivalent if either $x = y$ or $x, y \in X_n$ and the “tails” of x, y in $X_n = F_n \times C_{n+1} \times \cdots$ coincide starting from the second coordinate. If $x = (f_n, c_{n+1}(i_{n+1}), \dots) \in X_n$ then we let $\bar{x}^n := (f_n, 0, 0, \dots) \in X_n$. It now follows from (2-2) that

$$\alpha(x, T^{h_n}x) = \beta(\bar{x}^n, 0) \cdot a \quad \text{for all } x \in X'_n := \bigsqcup_{0 \leq i < i_n-1} [F_n + c_{n+1}(i)]_{n+1}.$$

For each $b \in \text{Orb}_K(a)$, we let

$$Y_{n,b} := \left\{ (x, k) \in X'_n \times K \mid k \cdot \alpha(x, T^{h_n}x) = b \right\}.$$

It is easy to see that

($\heartsuit$) $\{Y_{n,b} \mid b \in \text{Orb}_K(a)\}$ is a partition of $X'_n \times K$ into subsets of equal measure. Moreover,

$$\begin{aligned} (3-2) \quad & \sum_{j=1}^{i_n-1} \chi(k \cdot \alpha(x, T^{h_n}x))1_{[B+c_{n+1}(j-1)]_{n+1}}(x)\eta(k) \\ & = \sum_{b \in \text{Orb}_K(a)} \chi(b)1_{Y_{n,b}}(x, k)1_{[B]_n}(x)\eta(k) + \bar{o}(1). \end{aligned}$$

We claim that for each $b \in \text{Orb}_K(a)$,

($\diamond$) the set $Y_{n,b}$ is $\mathcal{R}_n(\beta)$ -invariant,

i.e. if $(x, k) \in Y_{n,b}$ and $(x, y) \in \mathcal{R}_n$ then $(y, k\beta(x, y)) \in Y_{n,b}$. This follows from the following fact:

$$\beta(\bar{x}^n, 0) = \beta(\bar{x}^n, \bar{y}^n)\beta(\bar{y}^n, 0) = \beta(x, y)\beta(\bar{y}^n, 0)$$

whenever $(x, y) \in \mathcal{R}_n$.

Since $\bigcup_{n>0} \mathcal{R}_n(\beta) = \mathcal{R}(\beta)$ and β is ergodic and the unit ball of bounded linear operators in $L^2(X \times K, \mu \times \lambda_K)$ is weakly compact and the algebra $L^\infty(X \times K)$ is weakly closed and $\mu(X'_n) \rightarrow \delta$, we deduce from $(\diamond)$ and $(\heartsuit)$ that $1_{Y_{n,b}}$ weakly converges to $\delta(\#\text{Orb}_K(a))^{-1}I$ for each $b \in \text{Orb}_K(a)$ as $\mathcal{N}_a - 1 \ni n \rightarrow \infty$. Hence

$$(3-3) \quad \sum_{b \in \text{Orb}_K(a)} \chi(b) 1_{Y_{n,b}} \rightarrow \delta l_\chi(a) I \quad \text{weakly as } \mathcal{N}_a - 1 \ni n \rightarrow \infty.$$

We now show that the cocycle $\tilde{\alpha}$ (or, equivalently, γ) is ergodic. Take arbitrary $n > 0$, $a \in A_0$, subset $B \subset F_n$ and an open subset $U \subset K$. Select $m > n$ such that $m \in \mathcal{N}_a - 1$ and

$$\mu \times \lambda_K(([B]_n \times U) \cap Y_{m,a}) > \frac{\delta}{2\#\text{Orb}_K(a)} (\mu \times \lambda_K)([B]_n \times U).$$

If $(x, k) \in ([B]_n \times U) \cap Y_{m,a}$ then $T_\beta^{h_m}(x, k) \in [B]_n \times U$ and

$$\tilde{\alpha}((x, k), T_\beta^{h_m}(x, k)) = k \cdot \alpha(x, T^{h_m}x) = a.$$

By the standard criterium of ergodicity for cocycles, the A_j -projection of $\tilde{\alpha}$ is ergodic for every $j > 0$. It follows that $\tilde{\alpha}$ (or γ) is ergodic.

Finally, we deduce (ii) from (3-1) by applying (3-3) and (3-2). To see that the second sum in (3-1) vanishes in the limit one should repeat the corresponding argument from the proof of (i) almost literally and make use of the ergodicity of γ (instead of β). $\square$

While proving the above proposition we established as a byproduct that γ is ergodic. We now show a stronger fact.

Corollary 3.5. *The γ -skew product extension T_γ of T is weakly mixing.*

Proof. If $T_\gamma v = \lambda v$ for a vector $0 \neq v \in L_0^2(X \times G, \mu \times \lambda_G)$ and some $\lambda \in \mathbb{T}$ then $|\langle U_{T_\gamma}^q v, v \rangle| = \|v\|^2$ for each $q > 0$. On the other hand, it follows from Proposition 3.4(i) and (ii) that $|\langle U_{T_\gamma}^{h_n} v, v \rangle| \rightarrow \delta \|v\|^2$ as $\mathcal{N}_0 - 1 \ni n \rightarrow \infty$, a contradiction. $\square$

Given $b \in A$ and $k \in K$, let L_b denote the operator of translation by b along the A -coordinate in $L^2(X \times K \times A, \mu \times \lambda_K \times \lambda_A)$ and let L_k denote the operator of translation by k along the K -coordinate in $L^2(X \times K, \mu \times \lambda_K)$.

Remark 3.6. The assertion of Proposition 3.4 can also be restated in the following equivalent way:

$$\begin{aligned} U_{T_\gamma}^{h_n} &\rightarrow \delta \cdot \frac{\sum_{b \in \text{Orb}_K(a)} L_b}{\#\text{Orb}_K(a)} + (1 - \delta)P_0 \quad \text{weakly as } \mathcal{N}_a - 1 \ni n \rightarrow \infty \quad \text{and} \\ U_{T_\beta}^{h_n} &\rightarrow \delta L_k + (1 - \delta)P_0 \quad \text{weakly as } \mathcal{L}_k - 1 \ni n \rightarrow \infty \end{aligned}$$

for each $a \in A_0$ and $k \in K_0$.

Given $\chi, \chi' \in \hat{A}$, we write $\chi \sim \chi'$ if there is $k \in K$ with $\chi = \chi' \circ k$.

Theorem 3.7. $\mathcal{M}((T_\beta)_{\tilde{\alpha}, H}) = L(K, \widehat{A}, \widehat{A/H}) = E$.

Proof. Since $U_{T_\gamma, H} = \bigoplus_{\chi \in \widehat{A/H}} U_{T_\beta, \chi}$, it is enough to show the following:

- (a) $U_{T_\beta, \chi}$ has a simple spectrum for each $\chi \in \widehat{A}$,
- (b) $U_{T_\beta, \chi}$ and $U_{T_\beta, \chi'}$ are unitarily equivalent if $\chi' \sim \chi$,
- (c) the measures of maximal spectral type of $U_{T_\beta, \chi}$ and $U_{T_\beta, \chi'}$ are mutually singular if $\chi' \not\sim \chi$.

We first note that (b) follows from (1-1).

Now, if $\chi' \neq \chi \circ k$ for any $k \in K$ then there is $a \in A_0$ such that $l_\chi(a) \neq l_{\chi'}(a)$. This follows from (ii) and (iii) of Lemma 3.2. Pass to the weak limits in the sequences $U_{T_\gamma, \chi}^{h_n}$ and $U_{T_\beta, \chi'}^{h_n}$ as $\mathcal{N}_a - 1 \ni n \rightarrow \infty$. Then (c) follows from (i) and (ii) of Proposition 3.4.

It remains to show (a). For this purpose we decompose $U_{T, \beta}$ into a direct orthogonal sum

$$U_{T_\beta} = \bigoplus_{\eta \in \widehat{K}} U_{T, \eta}.$$

Since T is of rank one and the function $X \ni x \mapsto \eta(\beta(x, Tx)) \in \mathbb{T}$ is constant on each cylinder $[f]_n$, $0 \leq f \leq h_n - 2$, it follows that the operator $U_{T, \eta}$ has a simple spectrum. We also deduce from (iii) and (iv) of Proposition 3.4 that the maximal spectral types of $U_{T, \eta}$ and $U_{T, \eta'}$ are mutually disjoint if $\eta \neq \eta'$. Hence U_{T_β} has a simple spectrum. Moreover, it is easy to verify that

- for each n , vectors $1_{[f]_n} \otimes 1_{\pi_n^{-1}(k)}$, $0 \leq f \leq h_n - 2$, $k \in K_n$, belong to the U_{T_β} -cyclic space generated by $\sum_{\eta \in \widehat{K}_n} 1_{[0]_n} \otimes \eta$ (at least up to small ϵ in the L^2 -norm) and
- the function

$$X \times K \ni (x, k) \mapsto (\chi(k \cdot \alpha(x, Tx)), \beta(x, Tx)) \in \mathbb{T}^2$$

is constant on each ‘level’ $[f]_n \times \pi_n^{-1}(k)$, $0 \leq f < h_n - 1$, $k \in K_n$.

These two facts imply (a). $\square$

We note that the condition (a) and (c) together imply

$$(3-5) \quad U_{T_\beta, \chi} \oplus U_{T_\beta, \xi} \text{ has a simple spectrum for all } \xi, \chi \in \widehat{A}, \chi \not\sim \xi.$$

In turn, (3-5) implies

- T_γ is weakly mixing and
- $\mathcal{M}(T_{\gamma, H}) = E$.

To force mixing we fix a sequence of positive reals δ_j that goes to 0 very fast and construct for each $j > 0$,

- a (C, F) -system $(X^{(j)}, \mu_j, T_j)$ associated to some sequence $(C_n^{(j)}, F_{n-1}^{(j)})_{n>0}$ and
- a cocycle $\gamma^{(j)} = (\beta^{(j)}, \alpha^{(j)}) : \mathcal{R}_j \rightarrow G = K \ltimes A$ associated to some sequence $(\beta_n^{(j)}, \alpha_n^{(j)})_{n>0}$

such that (3-5) is satisfied for the pair $(T_j, \gamma^{(j)})$ and the set $C_{n+1}^{(j)}$ is the union of an arithmetic sequence $\{0, h_n^{(j)}, \dots, (i_n^{(j)} - 1)h_n^{(j)}\}$ with a *staircase* sequence

$$\left\{ i_n^{(j)} h_n^{(j)} + 1, \dots, (r_n^{(j)} - 1)h_n^{(j)} + \frac{(r_n^{(j)} - i_n^{(j)})(r_n^{(j)} - i_n^{(j)} + 1)}{2} \right\}$$

and $i_n^{(j)}/r_n^{(j)} \rightarrow \delta_j$ as $n \rightarrow \infty$. Of course, $\mathcal{R}_j$ stands here for the T_j -orbit equivalence relation on $X^{(j)}$. In what follows we call such $(T_j, \gamma^{(j)})$ a δ_j -pair. The pair to be found will appear as a certain limit of the sequence $(T_j, \gamma^{(j)})$. Our main concern is to preserve (3-5) in the limit.

In the beginning of this section we outlined an explicit algorithm how to construct δ -pairs for an arbitrary $\delta > 0$, an arbitrary *initial* set F_0 and an arbitrary sequence $r_n \rightarrow \infty$ (provided that $\mu(X) < \infty$).

We carry out the construction of the sequence $(T_j, \gamma^{(j)})$ via an inductive procedure in j .

Let $(T_1, \gamma^{(1)})$ be an arbitrary δ_1 -pair. Suppose now that we have already built $(T_j, \gamma^{(j)})$. It follows from the inverse limit structure of K that

$$l^2(K_1) \subset l^2(K_2) \subset \dots$$

and $\bigcup_{m>0} l^2(K_m)$ is dense in $L^2(K, \lambda_K)$. Denote by $\mathcal{H}_s^{(j)}$ the finite dimensional linear space $\bigoplus_{f \in F_s^{(j)}} 1_{[f]_s} \otimes l^2(K_s)$. Then $\mathcal{H}_1^{(j)} \subset \mathcal{H}_2^{(j)} \subset \dots$ and $\bigcup_{s>0} \mathcal{H}_s^{(j)}$ is dense in $L^2(X_j \times K, \mu_j \times \lambda_K)$. Since (3-5) is satisfied for $(T_j, \gamma^{(j)})$, there exist $s_j > 0$, $Q_j > 0$ and a vector $v_j \in \mathcal{H}_{s_j}^{(j)} \oplus \mathcal{H}_{s_j}^{(j)}$ such that

- for each $w \in \mathcal{H}_{s_j}^{(j)} \oplus \mathcal{H}_{s_j}^{(j)}$,

$$\inf_{a_{-q}, \dots, a_q \in \mathbb{R}} \left\| w - \sum_{|q| \leq Q_j} a_q \left(U_{(T_j)_{\beta^{(j)}, \chi}}^q \oplus U_{(T_j)_{\beta^{(j)}, \xi}}^q \right) v_j \right\|_j \leq \delta_j \|w\|_j$$

whenever $\chi \not\sim \xi \in \widehat{A}_j$ and

- $U_{(T_j)_{\gamma^{(j)}, \chi}}^q v \in \mathcal{H}_{s_j}^{(j)}$ for all $|q| \leq Q$ and $\chi \in \widehat{A}_j$,

where $\|\cdot\|_j$ denotes the norm in $L^2(X^{(j)} \times K, \mu_j \times \lambda_K)$. Now let (T_{j+1}, γ_{j+1}) be any δ_{j+1} -pair whose initial set $F_0^{(j+1)}$ is $F_{s_j}^{(j)}$ and the *sequence of cuts* $(r_n^{(j+1)})_{n>0}$ is $j+1, j+2, \dots$. Thus, we now think that the sequence $(C_{n+1}^{(j+1)}, F_n^{(j+1)})_{n \geq 0}$ is constructed. It remains to repeat this process infinitely many times.

We now construct a certain “ (C, F) -limit” for the sequence $(T_j, \gamma^{(j)})_{j=1}^\infty$. For that we just let

$$\begin{aligned} s_0 &:= 0, \\ F_n &:= F_{n-s_1-\dots-s_{j-1}}^{(j)} & \text{if } s_0 + \dots + s_{j-1} \leq n < s_1 + \dots + s_j, \\ C_n &:= C_{n-s_1-\dots-s_{j-1}}^{(j)} & \text{if } s_0 + \dots + s_{j-1} < n \leq s_1 + \dots + s_j \quad \text{and} \\ \gamma_n &:= \gamma_{n-s_1-\dots-s_{j-1}}^{(j)} & \text{if } s_0 + \dots + s_{j-1} < n \leq s_1 + \dots + s_j. \end{aligned}$$

Denote by (X, μ, T) the (C, F) -system associated to $(C_n, F_{n-1})_{n>0}$. Let γ stand for the cocycle associated to $(\gamma_n)_{n>0}$. We now show that $T_{\gamma, H}$ provides a solution of Theorem 3.1.

Proof of Theorem 3.1. Since $\delta_j \rightarrow 0$, T is an almost staircase transformation. Moreover, $\#C_n \rightarrow \infty$ and $\#C_n \leq n$ for all $n \in \mathbb{N}$. Hence T is mixing [Ad]. Moreover, it is mixing of all orders since T is of rank one [Ka], [Ry1]. We also note that

$$F_{s_1+\dots+s_j} = F_0^{(j+1)} = F_{s_j}^{(j)}.$$

Consider a mapping Φ_j which sends a vector $1_{[f]_{s_j}} \otimes v \in L^2(X_j, \mu_j) \otimes l^2(K_{s_j})$ to a vector $1_{[f]_{s_1+\dots+s_j}} \otimes v \in L^2(X, \mu) \otimes l^2(K_{s_j})$, $f \in F_{s_j}^{(j)}$, $v \in l^2(K_{s_j})$. Then Φ_j extends naturally to a linear isomorphism of $\mathcal{H}_{s_j}^{(j)}$ onto the subspace

$$\mathcal{H}_{s_1+\dots+s_j} := \bigoplus_{f \in F_{s_1+\dots+s_j}} 1_{[f]_{s_1+\dots+s_j}} \otimes l^2(K_{s_j}) \subset L^2(X \times K, \mu \times \lambda_K)$$

and $\|\Phi_j v\| = \sqrt{\frac{\mu(X_{s_1+\dots+s_j})}{\mu_j(X_{s_j})}} \|v\|_j$ for all $v \in \mathcal{H}_{s_j}^{(j)}$. Moreover,

$$\Phi U_{(T_j)_{\beta(j)}, \chi}^q v = U_{T_{\beta}, \chi}^q \Phi v$$

whenever $v, U_{(T_j)_{\gamma(j)}, \chi}^q v \in \mathcal{H}_{s_j}^{(j)}$. Therefore for each $j > 0$, there exist $Q_j > 0$ and a vector $v_j \in \mathcal{H}_{s_1+\dots+s_j} \oplus \mathcal{H}_{s_1+\dots+s_j}$ such that for each $w \in \mathcal{H}_{s_1+\dots+s_{j-1}} \oplus \mathcal{H}_{s_1+\dots+s_{j-1}}$,

$$\inf_{a_{-q}, \dots, a_q \in \mathbb{R}} \left\| w - \sum_{|q| \leq Q_j} a_q \left(U_{T_{\gamma}, \chi}^q \oplus U_{T_{\gamma}, \xi}^q \right) v_j \right\| \leq \delta_j \|w\|$$

whenever $\chi \not\sim \xi \in \widehat{A}$. This yields (3-5). As we have already mentioned, (3-5) implies $\mathcal{M}(T_{\gamma, H}) = E$ and that T_{γ} is weakly mixing. Since T is mixing of all orders, it follows that T_{γ} is mixing of all orders [Ru]. Hence $T_{\gamma, H}$ is mixing of all orders because it is a factor of T . $\square$

4. MIXING TRANSFORMATIONS WHOSE SET OF SPECTRAL MULTIPLICITIES CONTAINS 2

Let E be an arbitrary subset of $\mathbb{N}$ such that $2 \in E$. Our main purpose in this section is to prove the following theorem.

Theorem 4.1. *There exists a mixing (of all orders) transformation S such that $\mathcal{M}(S) = E$.*

We first establish a generalization of the Algebraic lemma from [KL] (cf. with [Da5]).

Lemma 4.2. *Let P be a subset of $\mathbb{N}$. Then there is a triplet (K, B, D) such that $L(K, B, D) = P$. Moreover, if P is finite then we can choose B finite and K finite and cyclic.*

Proof. Let $P = \{p_1, p_2, \dots\}$ with $p_1 < p_2 < \dots$. For each $i > 0$, we select a finite group B_i and an automorphism θ_i of B_i such that for each $0 \neq b \in B_i$, the length of the θ_i -orbit of b is p_i . We now set

$$B := B_1 \oplus B_2^{\oplus p_1} \oplus B_3^{\oplus (p_1 p_2)} \dots,$$

$$D := \{(g_1, g_2, \dots) \in B \mid g_i = 0 \text{ if } i \notin \{1, 2, 2 + p_1, 2 + p_1 + p_1 p_2, \dots\}\}.$$

We consider two automorphisms of $B_i^{\oplus (p_1 \cdots p_{i-1})}$. The first one, say σ_i , is generated by the cyclic permutation of the coordinates. The second one, say θ'_i , is the Cartesian product $\theta_i \times \text{id} \times \dots \times \text{id}$. We note that $(\theta'_i \sigma_i)^{p_1 \cdots p_{i-1}} = \theta_i \times \dots \times \theta_i$. We now define an automorphism θ of B by setting

$$\theta := \theta_1 \times \theta'_2 \sigma_2 \times \theta'_3 \sigma_3 \times \dots.$$

We consider B equipped with θ as a $\mathbb{Z}$ -module. Let us compute the set $L(\mathbb{Z}, B, D)$. Take $0 \neq g = (g_1, g_2, \dots) \in D$. Let l be the maximal non-zero coordinate of g . If $l > 2$ then $l = 2 + \sum_{i=1}^k p_1 \cdots p_k$ for some $k > 0$. It is easy to see that

$$\min\{i > 1 \mid \text{such that } \theta^i g \in D\} = p_1 \cdots p_{k+1}.$$

Moreover, $\theta^{p_1 \cdots p_{k+1}} g = (g_1, \dots, g_{l-1}, \theta_{k+2} g_l, 0, 0, \dots)$. Hence the length of the θ -orbit of g is p_{k+2} . The remaining cases, when $l = 1, 2$, are considered in a similar way. We thus obtain $L(\mathbb{Z}, B, D) = P$. Passing to a compactification as in the proof of Lemma 3.1, we obtain a compact Abelian group K such that B is a K -module and $L(K, B, D) = P$. $\square$

From now on and till the end of the section A denotes the dual group $\widehat{B}$ and H denotes the dual group $\widehat{B/D} \subset A$. As follows from the proof of Lemma 4.2 we may assume without loss of generality that the conditions (i)–(iii) of Lemma 3.1 hold now as well. We restate and prove Theorem 4.1 in the following way.

Theorem 4.3. *There exists a mixing (of all orders) transformation S such that $\mathcal{M}(S) = L(K, \widehat{A}, \widehat{A/H}) \cup \{2\}$.*

We now state two auxiliary lemmata.

Lemma 4.4 ([Ag1], [Ry3]). *Let V be a unitary operator with a simple continuous spectrum.*

- (i) *If the weak closure of powers of V contains $\delta(I + V)$ for some $\delta > 0$ then $V \odot V$ has a simple spectrum.*
- (ii) *If $V \odot V$ has a simple spectrum then $V \otimes V$ has a homogeneous spectrum of multiplicity 2.*

Lemma 4.5 ([KaL]). *Let V_i , $i = 1, 2$, be unitary operators with a simple spectrum. Assume moreover that*

- (i) $V_i^{n_t} \rightarrow \delta(I + V_i^*)$ weakly, $i = 1, 2$, and
- (ii) $V_i^{m_t} \rightarrow \delta(I + c_i V_i^*)$ weakly, $i = 1, 2$

for some $\delta > 0$. If $c_1 \neq c_2$ then $V_1 \otimes V_2$ has also a simple spectrum.

Let $0 < i_n \leq r_n$, $r_n \rightarrow \infty$, $i_n/r_n \rightarrow \delta$, as $n \rightarrow \infty$, where $0 < \delta < 1$. Consider a partition

$$\mathbb{N} = \bigsqcup_{a \in A_0} (\mathcal{N}_a \sqcup \mathcal{M}_a) \sqcup \bigsqcup_{k \in K_0} \mathcal{L}_k$$

of $\mathbb{N}$ into infinite subsets $\mathcal{N}_a$, $\mathcal{M}_a$, $\mathcal{L}_k$.

If $n+1 \in \bigsqcup_{a \in A_0} \mathcal{N}_a \sqcup \bigsqcup_{k \in K} \mathcal{L}_k$ then we define $C_{n+1}, \alpha_{n+1}, \beta_{n+1}$ by exactly the same formulae as in Section 3. If $n+1 \in \mathcal{M}_a$ for some $a \in A_0$, we let

$$c_{n+1}(i) := \begin{cases} 0, & \text{if } i = 0, \\ c_{n+1}(i-1) + h_n, & \text{if } 0 < i < i_n, \\ c_{n+1}(i-1) + h_n + 1, & \text{if } i_n \leq i < 2i_n, \\ c_{n+1}(i-1) + h_n + i - 2i_n, & \text{if } 2i_n \leq i < r_n, \end{cases}$$

$$C_{n+1} := \{c_{n+1}(i) \mid 0 \leq i < r_n\},$$

$$\alpha_{n+1}(c_{n+1}(i)) := \begin{cases} 0, & \text{if } i = 0, \dots, i_n - 1, \\ \alpha_{n+1}(c_{n+1}(i-1)) - a, & \text{if } i = i_n, \dots, 2i_n - 1, \\ \alpha_{n+1}(c_{n+1}(i-1)), & \text{if } i = 2i_n, \dots, r_n - 1. \end{cases}$$

and $\beta_{n+1} \equiv 1$ on C_{n+1} .

Let (X, μ, T) stand for the (C, F) -system associated with $(C_{n+1}, F_n)_{n \geq 0}$. Denote by $\gamma = (\beta, \alpha) : \mathcal{R} \rightarrow G$ the (C, F) -cocycle associated with $(\beta_n, \alpha_n)_{n > 0}$.

Proposition 4.6. *For each $a \in A_0$,*

- (i) $U_{T_{\beta,1}}^{h_n} \rightarrow \delta I + (1 - \delta)P_0$ weakly as $\mathcal{N}_a - 1 \ni n \rightarrow \infty$, where P_0 is the orthogonal projection onto the subspace of constant functions on $X \times K$.
- (ii) $U_{T_{\beta,\chi}}^{h_n} \rightarrow \delta l_\chi(a)I$ weakly as $\mathcal{N}_a - 1 \ni n \rightarrow \infty$ for each $1 \neq \chi \in \hat{A}$.
- (iii) $U_{T_{\beta,1}}^{h_n} \rightarrow \delta(I + U_{T_{\gamma,1}}^*) + (1 - 2\delta)P_0$ weakly as $\mathcal{M}_a - 1 \ni n \rightarrow \infty$, where P_0 is the orthogonal projection onto the subspace of constant functions on $X \times K$.
- (iv) $U_{T_{\beta,\chi}}^{h_n} \rightarrow \delta(I + l_\chi(a)U_{T_{\gamma,\chi}}^*)$ weakly as $\mathcal{M}_a - 1 \ni n \rightarrow \infty$ for each $1 \neq \chi \in \hat{A}$.

For each $k \in K$,

- (v) $U_{T,1}^{h_n} \rightarrow \delta I + (1 - \delta)P_0$ weakly as $\mathcal{L}_k - 1 \ni n \rightarrow \infty$, where P_0 is the orthogonal projection onto the subspace of constant functions on X .
- (vi) $U_{T,\eta}^{h_n} \rightarrow \delta \eta(k)I$ weakly as $\mathcal{L}_k - 1 \ni n \rightarrow \infty$ for each $1 \neq \eta \in \hat{K}$.

Proof. Repeating the argument from the proof of Proposition 3.2 verbally we obtain (i), (ii), (v), (vi). The remaining claims can be shown in a similar way. For instance, take $n \in \mathcal{M}_a - 1$, subsets $B \subset F_n$ and $\eta \in \hat{K}$. Then we obtain three “sums” (instead

of the two sums in the proof of Proposition 3.2) in the righthandside of the following:

$$\begin{aligned}
(U_{T,\eta}^{h_n} 1_{[B]_n})(x) &= \eta(\beta(x, T^{h_n} x)) 1_{T^{-h_n}[B]_n}(x) \\
&= \sum_{j=1}^{i_n-1} 1_{[B+c_{n+1}(j-1)]_{n+1}}(x) \\
&\quad + \sum_{j=i_n}^{2i_n-1} U_{T,\eta}^* 1_{[B+c_{n+1}(j-1)]_{n+1}}(x) \\
&\quad + \sum_{j=2i_n+1}^{r_n-1} U_{T,\eta}^{2i_n-j} 1_{[B+c_{n+1}(j-1)]_{n+1}}(x) + \bar{o}(1)
\end{aligned}$$

for all $x \in X$ and $k \in K$. Hence

$$U_{T,\eta}^{h_n} \rightarrow \delta(I + U_{T,\eta}^*) + (1 - 2\delta)P_\eta \text{ weakly as } \mathcal{M}_a - 1 \ni n \rightarrow \infty.$$

The claim (iii) follows. Similar minor “changes” of the proof of Proposition 3.2 are needed to establish (iv). We leave details to the readers. $\square$

We now prove a *partially rigid* version of Theorem 4.3.

Theorem 4.7. $\mathcal{M}(T_{\gamma,H} \times T_\beta) = L(K, \widehat{A}, \widehat{A/H}) \cup \{2\}$.

Proof. Since $U_{T_{\gamma,H}} = \bigoplus_{\chi \in \widehat{A/H}} U_{T_\beta,\chi}$, it is enough to show the following:

- (a) $U_{T_\beta,\chi} \otimes U_{T_\beta}$ has a simple spectrum for each $1 \neq \chi \in \widehat{A}$,
- (b) $U_{T_\beta} \otimes U_{T_\beta}$ has a homogeneous spectrum of multiplicity 2 (in the orthocomplement to the constants),
- (c) $U_{T_\beta,\chi} \otimes U_{T_\beta}$ and $U_{T_\beta,\chi'} \otimes U_{T_\beta}$ are unitarily equivalent if $\chi' \sim \chi$,
- (d) the measures of maximal spectral type of $U_{T_\beta,\chi} \otimes U_{T_\beta}$ and $U_{T_\beta,\chi'} \otimes U_{T_\beta}$ are mutually singular if $\chi' \not\sim \chi$.

It follows from (v) and (vi) of Proposition 4.6 that $U_{T_\beta,\chi}$ has a simple spectrum for each $\chi \in \widehat{A}$ (it was shown in the proof of Theorem 3.7). Denote by $U_{T_\beta}^\circ$ the restriction of U_{T_β} to the orthocomplement to the constants. Then making use of (iii) and (iv) of Proposition 4.6 we deduce from Lemma 4.5 that $U_{T_\beta,\chi} \otimes U_{T_\beta}^\circ$ has a simple spectrum if $\chi \neq 1$. On the other hand, it follows from (i) and (ii) of Proposition 4.6 that

$$(4-1) \quad (U_{T_\beta,\chi} \otimes U_{T_\beta}^\circ)^{h_n} \rightarrow \delta^2 I \text{ and } U_{T_\beta,\chi}^{h_n} \rightarrow \delta I \text{ as } \mathcal{N}_0 - 1 \ni n \rightarrow \infty.$$

Therefore the maximal spectral types of $U_{T_\beta,\chi} \otimes U_T^\circ$ and $U_{T_\beta,\chi}$ are mutually disjoint. Hence the orthogonal sum of these two operators, i.e. $U_{T_\beta,\chi} \otimes U_T$, has a simple spectrum. Thus (a) is proved.

We deduce from Lemma 4.4 and Proposition 4.6(iii) that $U_{T_\beta}^\circ \otimes U_{T_\beta}^\circ$ has a homogeneous spectrum of multiplicity 2. In view of Proposition 4.6(i),

$$(U_{T_\beta}^\circ \otimes U_{T_\beta}^\circ)^{h_n} \rightarrow \delta^2 I \text{ and } (U_{T_\beta}^\circ)^{h_n} \rightarrow \delta I \text{ as } \mathcal{N}_0 - 1 \ni n \rightarrow \infty,$$

it follows that $U_{T_\beta}^\circ \otimes U_{T_\beta}^\circ$ and $U_{T_\beta}^\circ$ are spectrally disjoint. Therefore we obtain (b).

(c) follows from (1-1).

To show (d), we first find $a \in A_0$ such that $l_\chi(a) \neq l_{\chi'}(a)$. It follows from (i) and (ii) of Proposition 4.6 that

$$(U_{T_\beta, \chi} \otimes U_{T_\beta}^\circ)^{h_n} \rightarrow \delta^2 l_\chi(a) I \quad \text{and} \quad (U_{T_\beta, \chi'} \otimes U_{T_\beta}^\circ)^{h_n} \rightarrow \delta^2 l_{\chi'}(a) I \quad \text{as } \mathcal{N}_a - 1 \ni n \rightarrow \infty.$$

Hence $U_{T_\beta, \chi} \otimes U_{T_\beta}^\circ$ and $U_{T_\beta, \chi'} \otimes U_{T_\beta}^\circ$ are spectrally disjoint. From (4-1) we deduce that $U_{T_\beta, \chi} \otimes U_{T_\beta}^\circ$ and $U_{T_\beta, \chi'}$ are spectrally disjoint as well as $U_{T_\beta, \chi'} \otimes U_{T_\beta}^\circ$ and $U_{T_\beta, \chi}$ are. Hence (d) follows. $\square$

Proof of Theorem 4.3. We note that the condition (a) and (d) from the proof of Theorem 4.7 together are equivalent to the following

$$(4-2) \quad (U_{T_\beta, \chi} \otimes U_{T_\beta}) \oplus (U_{T_\beta, \xi} \otimes U_{T_\beta}) \text{ has a simple spectrum for all } \chi \not\sim \xi \in \widehat{A}.$$

In view of Lemma 4.4(ii), the condition (b) of Theorem 4.7 is equivalent to

$$(4-3) \quad U_{T_\beta} \oplus (U_{T_\beta}^\circ \odot U_{T_\beta}^\circ) \text{ has a simple spectrum.}$$

Thus (4-2) and (4-3) yield that $\mathcal{M}(T_{\gamma, H} \times T_\beta) = L(K, \widehat{A}, \widehat{A/H}) \cup \{2\}$ and T_γ is weakly mixing.

It remains to force mixing in the same way as we did in the proof of Theorem 3.1. For that we need to construct an *approximating* sequence $(T_j, \gamma^{(j)})_{j>0}$ of δ_j -pairs with $\delta_j \rightarrow 0$ and pass to the (C, F) -limit. Our main concern now is to preserve (4-2) and (4-3) in the limit. This requires only a slight modification of the argument in the proof of Theorem 3.3. We leave details to the reader. $\square$

5. CONCLUDING REMARKS

- (1) Modifying slightly the techniques developed in this paper one can obtain other series of sets admitting mixing realizations. For instance, consider the spectral multiplicities of the Cartesian products $T_{\gamma, H} \times T_\beta^{\times r}$ with $r > 1$ and their natural factors.
- (2) Other mixing realizations may appear as Poissonian (and Gaussian) ones. For instance, given any $q > 2$, the geometric sequence $\{q, q^2, q^3, \dots\}$ admits a mixing Poissonian realization [DaR2].
- (3) We do not know, however, are there mixing realizations for $n \cdot E$, where $1 \in E \subset \mathbb{N}$ and $n > 2$. Weakly mixing realizations for such subsets were constructed in [Da3].

REFERENCES

- [Ad] T. M. Adams, *Smorodinsky's conjecture on rank one systems*, Proc. Amer. Math. Soc. **126** (1998), 739–744.
- [Ag1] O. N. Ageev, *On ergodic transformations with homogeneous spectrum*, J. Dynam. Control Systems **5** (1999), 149–152.
- [Ag2] ———, *On the multiplicity function of generic group extensions with continuous spectrum*, Ergodic Th. & Dynam. Systems **21** (2001), 321–338.
- [Ag3] ———, *The homogeneous spectrum problem in ergodic theory*, Invent. Math. **160** (2005), 417–446.
- [Ag4] ———, *Mixing with staircase multiplicity function*, Ergodic Th. & Dynam. Systems **28** (2008), 1687–1700.

- [Da1] A. I. Danilenko, *On cocycles with values in group extensions. Generic results*, Matemat. Fizika, Analiz, Geometriya **7** (2000), 153–171.
- [Da2] A. I. Danilenko, *Funny rank one weak mixing for nonsingular Abelian actions*, Isr. J. Math. **121** (2001), 29–54.
- [Da3] ———, *Explicit solution of Rokhlin’s problem on homogeneous spectrum and applications*, Ergod. Th. & Dyn. Syst. **26** (2006), 1467–1490.
- [Da4] ———, *(C, F)-actions in ergodic theory*, Geometry and Dynamics of Groups and Spaces, Progr. Math., vol. 265, 2008, pp. 325–351.
- [Da5] ———, *On new spectral multiplicities for ergodic maps*, preprint, arXiv:0904.4808.
- [DaR1] A. I. Danilenko and V. V. Ryzhikov, *Spectral multiplicities for infinite measure preserving transformations*, preprint, arXiv:0905.3486.
- [DaR2] A. I. Danilenko and V. V. Ryzhikov, *Mixing constructions with infinite invariant measure and spectral multiplicities*, preprint.
- [FM] J. Feldman and C. C. Moore, *Ergodic equivalence relations, cohomology, and von Neumann algebras. I*, Trans. Amer. Math. Soc. **234** (1977), 289–324.
- [G–Li] G. R. Goodson, J. Kwiatkowski, M. Lemańczyk and P. Liardet, *On the multiplicity function of ergodic group extensions of rotations*, Studia Math. **102** (1992), 157–174.
- [Kal] S. A. Kalikow, *Twofold mixing implies threefold mixing for rank one transformations*, Ergodic Theory & Dynam. Systems **4** (1984), 237–259.
- [Ka] A. B. Katok, *Combinatorial constructions in ergodic theory and dynamics*, University Lecture Series, 30, Amer. Math. Soc., Providence, RI, 2003.
- [KaL] A. Katok and M. Lemańczyk, *Some new cases of realization of spectral multiplicity function for ergodic transformations*, Fund. Math. (to appear).
- [KL] J. Kwiatkowski (jr) and M. Lemańczyk, *On the multiplicity function of ergodic group extensions. II*, Studia Math. **116** (1995), 207–215.
- [Ju] A. del Junco, *A simple map with no prime factors*, Israel J. Math. **104** (1998), 301–320.
- [Le] M. Lemańczyk, *Spectral theory of dynamical systems*, Encyclopedia of complexity and systems science, Springer, 2009.
- [Ne] D. Newton, *On Gaussian processes with simple spectrum*, Z. Wahrscheinlichkeitstheorie Werv. Gebiete **5** (1966), 207–209.
- [Os] V. I. Oseledec, *On the spectrum of ergodic automorphisms*, Soviet Math. Dokl. **168** (1966), 776–779.
- [Ro1] E. A. Robinson, *Ergodic measure-preserving transformations with arbitrary finite spectral multiplicities*, Invent. Math. **72** (1983), 299–314.
- [Ro2] E. A. Robinson, *Mixing and spectral multiplicity*, Erg. Theory & Dynam. Systems **5** (1985), 617–624.
- [Ro3] ———, *Transformations with highly nonhomogeneous spectrum of finite multiplicity*, Israel J. Math. **56** (1986), 75–88.
- [Ry1] V. V. Ryzhikov, *Mixing, rank and minimal self-joining of actions with invariant measure*, Mat. Sb. **183** (1992), 133–160.
- [Ry2] ———, *Transformations having homogeneous spectra*, J. Dynam. Control Systems **5** (1999), 145–148.
- [Ry3] ———, *Homogeneous spectrum, disjointness of convolutions, and mixing properties of dynamical systems*, Selected Russian Mathematics **1** (1999), 13–24.
- [Ry4] ———, *Weak limits of powers, the simple spectrum of symmetric products and mixing constructions of rank 1*, Sb. Math. **198** (2007), 733–754.
- [Ry5] ———, *Spectral multiplicities and asymptotic operator properties of actions with invariant measure*, Sb. Math. (to appear).
- [Ru] D. Rudolph, *k-fold mixing lifts to weakly mixing isometric extensions*, Ergodic Theory Dynam. Systems **5** (1985), 445–447.
- [Sc] K. Schmidt, *Cocycles of ergodic transformation groups*, Lecture Notes in Math. vol 1, McMillan Co of India, 1977.
- [Ti] S. V. Tikhonov, *A complete metric on the set of mixing transformations*, Sb. Math. **198** (2007), 575–596.